













CORRECTORS

BIOVAL Mn-Zn CITRICS

DEFICIENCIES CORRECTOR MANGANESE AND ZINC FORMULATED IN FLOW SPECIAL CITRUS



DECLARED CONTENT	
Zinc (Zn) Total	13,50% p/v
Manganese (Mn) Total	13,50% p/v
Nitrogen (N) Total	5,80% p/v

PHYSICOCHEMICAL CHARACTERISTICS

Density: 1,65

pH (solution 10%): 5,5-6,5



Its special formulation and high concentration helps to prevent and simultaneously correct deficiencies of manganese and zinc. The combined application of Zinc and Manganese is more effective than the application of each element separately. The low levels of zinc reduce the number of fruits per tree and, by extension, the size of the fruits. A symptom of zinc deficiency in citrus is the appearance of chlorotic leaf spots and / or whitish areas with green veins. On the other hand

whitish areas with green veins. On the other hand, manganese deficiency is frequently manifested by a speckle of leaves.

Thanks to the composition of **BIOVAL Mn-Zn CITRICS** it is possible to optimize the contribution of nutrients (Zn and Mn) in a more persistent way in the leaf, so that the period of effectiveness of the application is longer. With **BIOVAL Mn-Zn CITRICS** you will increase the size of the leaf, shoots and fruits, improve the vitamin C content and get more fruits per tree.





RECOMMENDED GENERAL DOSE:

FOLIAR: 250-450 cc / hl (2-4 applications during the cycle according to spring-summer needs).

Irrigation Drip: 4-10L / hectare (2-4 applications per cycle according to needs).

It is recommended, however, to follow the doses and applications established by their technicians at the field level, which will vary according to the type of crop, their extractions in fertilizer units and the phenological state of the same. From "Distribución del Levante Bioval, S.L." or through our agents or distributors we can advise you on the dose of use according to your crops.

STORAGE:

Look for storage in a dry place, cool and without long exposure to the sun. Keep in original container.

Shake before using

Away from food, drink and feed.

Do not stack more than four heights.

It is recommended to use it within 2 years of the lot date.

Precautions, risk phrases and safety: see label on packaging.



