

# MACRO



# Nutrients



CERO  
residuo



bival  
Distribución del Levante BIOVAL S. L

Linea

## MACRO-GEL

### BIOVAL DENSO SLOW N 44

NITROGEN WITH DCD

SLOW RELEASE



### DECLARED CONTENT

Total Nitrogen (N)	44,0% p/v
Nitrogen (N) Ammonia	10,56% p/v
Nitrogen (N) Nitric	10,56% p/v
Nitrogen (N) Amide	22,88% p/v
DCD (Dianandiamide)	0,30% p/v

### PHYSICO-CHEMICAL CHARACTERISTICS

Formulation: GEL

pH (solution 1%): 6-7

E.C. mS/cm (solution 1%): 0.8

Yellow color

Density (gr/cc): 1,32



### PROPERTIES

Fertilizer of high nitrogen content with dicyandiamide (DCD) inhibitor of nitrification to delay the transformation of ammoniacal and amidic nitrogen into nitric, a process that guarantees an assimilable and constant source of nitric nitrogen and the reduction to minimum losses of nitrogen by leaching. BIOVAL DENSO SLOW N 44 (DCD) manifests itself as a broad-spectrum fertilizer, rapid assimilation and economic use due to the homogeneity, precision and low dose required in its use (ammonium + ammoniac + nitric). Its use favors the growth and vegetative development, with vitalizing, stimulating and favorable effect of the absorption of macro and micronutrients and its effect on the increase of the production.



#### RECOMMENDED GENERAL DOSE:

Foliar: 150-300 cc/hl in 4-7 applications, according to culture.


Fertigation: 15-20 L/hectare and year (4-7 applications at a rate of 100-200 cc/hl) according to culture.

*It is recommended, however, to follow the doses and applications established by their technicians at the field level, which will vary according to the type of crop, their extractions in fertilizer units and the phenological state of the same. From "Distribución del Levante Bioval, S.L." or through our agents or distributors we can advise you as to the use dose according to your culture.*

#### STORAGE:

Stable product if not previously diluted.  
Store in the original package  
Look for storage in a dry place, cool and without long exposure to the sun.  
Away from food, drink and feed.  
Do not stack more than four heights.  
It is recommended to use it within 2 years of the lot date.  
Precautions, risk phrases and safety: see label on packaging.

#### PRESENTATION:



**Warnings:** For pivot, spray and drip, use suitable material pipes and use water alone at the end of irrigation for circuit washing.  
Avoid excessive doses as it may delay maturation and sensitize the fungal attack.  
Do not mix with sulfo-calcium broth, dinocap, sulfur or alkaline reaction products.  
Avoid treating during the hours of maximum heat.